

Comment on “Quasisaddles as relevant points of the potential energy surface in the dynamics of supercooled liquids” [J. Chem. Phys. 116, 10297 (2002); cond-mat/0203301]

Jonathan P. K. Doye, and David J. Wales
University Chemical Laboratory, Lensfield Road, Cambridge CB2 1EW, United Kingdom

Following theoretical work that emphasized the importance of saddle points on a potential energy surface (PES) in the dynamics of supercooled liquids,¹ two papers appeared proposing a mapping of instantaneous configurations onto the underlying saddle points of the PES.^{2,3} In this mapping the instantaneous configurations are starting points for minimizations on the function $W = |\nabla V|^2$, where V is the potential energy. The motivation for this choice was that the stationary points of the PES, of whatever index, correspond to absolute minima of W with $W = 0$. However, we subsequently pointed out that the vast majority of the minima of W do not correspond to stationary points of the PES, but have $W > 0$.⁴ This result was recently confirmed by the authors of Ref. 2 after reanalysing their data.⁵

The minima of W must satisfy $\nabla W = 2\mathbf{H}\mathbf{g} = \mathbf{0}$ where $\mathbf{H}$ is the Hessian (the second derivative matrix of the potential energy) and $\mathbf{g}$ is the gradient, ∇V . There are two types of solution to this equation. First, those where $\mathbf{g} = \mathbf{0}$, which correspond to true stationary points of the PES, and second, those where $\mathbf{g}$ is an eigenvector of the Hessian with zero eigenvalue. These minima have $W > 0$ and have a point of inflection in the direction of the gradient on the PES, and we will continue to refer to them as non-stationary points (NSP's) in order to differentiate them from the first set of minima.

In Ref. 5 Angelani *et al.* now argue that it is reasonable to consider the NSP's as ‘quasi-saddles’: ‘If this number [the number of inflection directions at the minima of W] is small one can think of the minima of W as a “quasi” saddle point (QSP), as they are *true* saddle points in the subspace orthogonal to the small number of inflection directions. One can conjecture that the properties of a QSP are very similar to that of a true saddle point associate (sic) to an instantaneous configuration’. In this comment we suggest that these properties simply follow from the form of the solutions to $\nabla W = 0$, and do not really indicate any saddle-like character.

We first note that NSP's are expected to have exactly one zero Hessian eigenvalue (in addition to the three zeros corresponding to translation of the whole system), unless there is some symmetry present in the system giving rise to degenerate eigenvalues, or an unexpected accidental degeneracy, both of which are very unlikely in the present case. This analytic result is confirmed by an anal-

ysis of the 8,000 stationary points of W of the 256-atom system in our database.⁴ Only eight of the 6,920 NSP's have a non-zero Hessian eigenvalue with magnitude less than 0.001. Using the convergence criteria described in Ref. 4 the average magnitude of the zero eigenvalue for the NSP's is 6×10^{-5} , and the average magnitude of the smallest non-zero eigenvalue over all 8,000 points is 0.64.⁶ We therefore suggest that the results in Ref. 5, where the average number of directions with zero curvature varies from between 1.1 and 4.4 (depending on temperature), are due to incomplete minimization, and that the tolerance used to assign the zero eigenvalues in Ref. 5 is not tight enough. Instead, the number of extra zero Hessian eigenvalues should be precisely one, independent of temperature.

Secondly, we note that at an NSP the inflection direction is parallel to the gradient. Hence the gradient in the sub-space orthogonal to the inflection direction is zero, and the NSP is a stationary point in this subspace. However, any point in configuration space is a stationary point in the subspace orthogonal to the gradient, and so this result does not imply any additional saddle-like properties.

The authors of Ref. 5 introduce a further justification for the use of the term ‘quasisaddle’ in a footnote: ‘most of the QSP found are “almost” saddles in the sense that W is almost zero numerically’. It is not surprising that the magnitude of the gradient at the NSP's is significantly smaller than at the instantaneous configuration, since this is what is being minimized. However, analysing our database shows that W is not almost zero numerically. The average value of W at a NSP varies between 0.28 and 3.25 depending on temperature, compared to a value for the true stationary points of 2×10^{-8} , i.e. numerically zero consistent with our convergence criteria.

We therefore disagree with the arguments used in Ref. 5 to infer saddle-like properties for NSP's. The most interesting results of Angelani *et al.* do not depend upon this classification—the observation that the number of negative eigenvalues of the NSP's extrapolates to zero at the mode-coupling temperature, and is strongly correlated with the magnitude of the diffusion coefficient, is still intriguing. However, the interpretation of these results depends upon a correct understanding of the nature of the NSP's.

¹ A. Cavagna, Europhys. Lett. **53**, 490 (2001).

² L. Angelani *et al.*, Phys. Rev. Lett. **85**, 5356 (2000).

³ K. Broderix *et al.*, Phys. Rev. Lett. **85**, 5360 (2000).

⁴ J. P. K. Doye and D. J. Wales, J. Chem. Phys. **116**, 3777 (2002).

⁵ L. Angelani *et al.*, J. Chem. Phys. **116**, 10297 (2002).

⁶ As in Ref. 4 we use ϵ_{AA} and σ_{AA} as the units of energy and distance.